

STRANGE CIRCLES

THE RILEY SLICE OF QUASI-FUCHSIAN SPACE

ALEX ELZENAAR


(MPI-MIS)

JOINT WORK WITH:
GAVEN MARTIN

(NZ INST. OF ADV. STUDY, MASSEY UNI., AUCKLAND)


JEROEN SCHILLEWAERT

(THE UNIVERSITY OF AUCKLAND)




APRIL 27, 2022

GEOMETRIC GROUP THEORY



GEOMETRIC GROUP THEORY



- A discrete subgroup of $\mathrm{PSL}(2, \mathbb{C})$ is called **Kleinian**.


- A discrete subgroup of $\mathrm{PSL}(2, \mathbb{C})$ is called **Kleinian**.
- Kleinian groups act as isometries of \mathbb{H}^3 , and as conformal maps on the ‘sphere at infinity’, $\hat{\mathbb{C}} := \mathbb{C} \cup \infty = \mathbb{P}\mathbb{C}^1$.

- A discrete subgroup of $\mathrm{PSL}(2, \mathbb{C})$ is called **Kleinian**.
- Kleinian groups act as isometries of \mathbb{H}^3 , and as conformal maps on the 'sphere at infinity', $\hat{\mathbb{C}} := \mathbb{C} \cup \infty = \mathbb{P}\mathbb{C}^1$.
- Discreteness is enough to make sure that \mathbb{H}^3/G is an orbifold.

- A discrete subgroup of $\mathrm{PSL}(2, \mathbb{C})$ is called **Kleinian**.
- Kleinian groups act as isometries of \mathbb{H}^3 , and as conformal maps on the 'sphere at infinity', $\hat{\mathbb{C}} := \mathbb{C} \cup \infty = \mathbb{P}\mathbb{C}^1$.
- Discreteness is enough to make sure that \mathbb{H}^3/G is an orbifold.
- It is **not** enough for $\hat{\mathbb{C}}/G$ to be a Riemann surface.

LIMIT SETS

Discrete subgroups of $PSL(2, \mathbb{C})$ have orbits which accumulate at 'limit points'.




Orbit of 0 under $\begin{bmatrix} 0.9 + 0.05i & 0.1 + 0.45i \\ 0.1 + 0.45i & 0.9 + 0.05i \end{bmatrix}$.

Source at -1 , sink at 1 ; orbit lies on a log-spiral.

AN EXAMPLE

$$\Gamma_{1+2i} = \left\langle \left[\begin{array}{cc} 1 & 1 \\ 0 & 1 \end{array} \right], \left[\begin{array}{cc} 1 & 0 \\ 1+2i & 1 \end{array} \right] \right\rangle$$



AN EXAMPLE, CONT.

$$\Gamma_{1+2i} = \left\langle \left[\begin{array}{cc} 1 & 1 \\ 0 & 1 \end{array} \right], \left[\begin{array}{cc} 1 & 0 \\ 1+2i & 1 \end{array} \right] \right\rangle$$

Theorem

Γ_{1+2i} is discrete and free on these two generators.

Theorem

The Riemann surface coming from Γ_{1+2i} is a 4-times punctured sphere.

These results are not supposed to be obvious!!!

THE MODULI SPACE ASSOCIATED TO THE EXAMPLE

Question

What happens when you wiggle the parameter $1 + 2i$ around in \mathbb{C} ?

'Wiggle' means 'move holomorphically'.

Theorem (Extended λ -lemma: Mañé, Sad, Sullivan (1983); Ślodkowski (1991); Earle, Kra, Krushkal' (1994))

*If a matrix coefficient in a generator of a Kleinian group moves holomorphically, then the limit set deforms **quasiconformally** until it collides with itself.*

If Γ_ρ is a group with parameter ρ , then $\Gamma_{\tilde{\rho}} = f^{-1}\Gamma_\rho f$ for some quasiconformal map $f : \hat{\mathbb{C}} \rightarrow \hat{\mathbb{C}}$.

The Riley slice is the quasiconformal moduli space of the example.

Definition

The **Riley slice** \mathcal{R} is the set of $\rho \in \mathbb{C}$ such that

$$\Gamma_\rho := \left\langle \begin{bmatrix} 1 & 1 \\ 0 & 1 \end{bmatrix}, \begin{bmatrix} 1 & 0 \\ \rho & 1 \end{bmatrix} \right\rangle \leq \mathrm{PSL}(2, \mathbb{C})$$


is quasiconformally conjugate to Γ_{1+2i} .

THE RILEY SLICE

The **Riley slice** \mathcal{R} is the set of $\rho \in \mathbb{C}$ such that

$$\Gamma_\rho := \left\langle \begin{bmatrix} 1 & 1 \\ 0 & 1 \end{bmatrix}, \begin{bmatrix} 1 & 0 \\ \rho & 1 \end{bmatrix} \right\rangle \leq \mathrm{PSL}(2, \mathbb{C})$$

is quasiconformally conjugate to Γ_{1+2i} .



BIG THEOREM 1

Theorem (Lyubich, Suvorov (1988); Maskit, Swarup (1989); Ohshika, Miyachi (2010); Aigner (conj. 2001); Akiyoshi, Ohshika, Parker, Sakuma, Yoshima (2020); Aimi, Lee, Sakai, Sakuma (2020))

The Riley slice \mathcal{R} is equivalently:

1. *The set of $\rho \in \mathbb{C}$ such that*

$$\Gamma_\rho := \left\langle \begin{bmatrix} 1 & 1 \\ 0 & 1 \end{bmatrix}, \begin{bmatrix} 1 & 0 \\ \rho & 1 \end{bmatrix} \right\rangle \leq \mathrm{PSL}(2, \mathbb{C})$$

is free and discrete, and such that the associated Riemann surface is homeomorphic to a 4-times punctured sphere.

2. *The interior of the set of ρ such that Γ_ρ is free and discrete.*

Proof is extremely non-trivial (especially part 2).

BIG THEOREM 2

Theorem (Lyubich, Suvorov (1988); Maskit, Swarup (1989); application of Bers, Greenberg, Marden (1971))

The Riley slice is homeomorphic to an annulus. More precisely,

$$\mathcal{R} \simeq \frac{\mathbb{H}^2}{\langle \omega \rangle}$$

where \mathbb{H}^2 is the Teichmüller space of 4-times punctured spheres and where $\omega \in \text{Mod}(S_{0,4})$ is explicit enough that the quotient is 'obviously' an annulus.

Proof is an exercise in applying Teichmüller theory.

BIG THEOREM 3

Theorem (Keen, Series (1994); Komori, Series (1998); generalised E., Martin, Schillewaert (2021-22))

The Riley slice is homeomorphic to an annulus. More precisely,

$$\mathcal{R} \simeq S^1 \times \mathbb{R}_{>0}$$

where the first coordinate chooses a distinguished geodesic on the Riemann surface, and where the second coordinate gives its measure.

This is a concrete case of the Ahlfors-Bers-Maskit deformation/parabolicity theory for Kleinian groups.

WHAT ABOUT THE 'ALGEBRAIC' MODULI SPACE OF 4-PUNCTURED SPHERES?

$$\begin{array}{ccc} \text{Teich}(S_{0,4}) \cong \mathbb{H}^2 & & \\ \downarrow \text{Mod}(S_{0,4}) & \searrow \langle \omega \rangle & \\ \mathcal{M}_{0,4} \cong S_{0,3} & & \mathcal{R} \cong \mathbb{R}_{>0} \times \mathbb{R}/2\mathbb{Z} \end{array}$$

↙ ?

WHAT ABOUT THE 'ALGEBRAIC' MODULI SPACE OF 4-PUNCTURED SPHERES?

$$\begin{array}{ccc} \text{Teich}(S_{0,4}) \cong \mathbb{H}^2 & & \\ \downarrow \text{Mod}(S_{0,4}) & \searrow \langle \omega \rangle & \\ & & \mathcal{R} \cong \mathbb{R}_{>0} \times \mathbb{R}/2\mathbb{Z} \\ & \swarrow \text{knot theory} & \\ \mathcal{M}_{0,4} \cong S_{0,3} & & \end{array}$$

BEDTIME READING

- Very introductory survey for people in other fields: A.J.E., G.M., J.S. “Concrete one complex dimensional moduli spaces of hyperbolic manifolds and orbifolds”. In: *2021-22 MATRIX annals* (Springer, to appear; arXiv 2022).
- More detailed introduction: A.J.E. *Deformation spaces of Kleinian groups*. MSc thesis. The Uni. of Auckland (2022).
- Our recent work: A.J.E., G.M., J.S. “Approximations of the Riley slice” (arXiv 2021), “The combinatorics of Farey words and their traces” (arXiv 2022), “The Riley slice and its elliptic cousins” (to appear soon).
- Title picture: Hilma af Klint, *The SUW series, group IX*, 1915, oil on canvas. City Gallery Wellington (2021–2022), exhibition courtesy of the Hilma af Klint Foundation. Photo: Cheska Brown.